

AD-A072 400

MARYLAND UNIV BALTIMORE SCHOOL OF MEDICINE
HEMOGLOBIN FUNCTION IN STORED BLOOD.(U)
JAN 79 R B DAWSON

F/G 6/5


UNCLASSIFIED

DADA17-72-C-2005
NL

| OF |
AD
A072400



END
DATE
FILMED
9-79
DDC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

LEVEL

4

HEMOGLOBIN FUNCTION IN STORED BLOOD

Annual Report

January 1979

(for the period 1 February 1978 - 31 January 1979)

by

R. Ben Dawson, M. D.

Supported by


US Army Medical Research and Development Command
Fort Detrick, Frederick, Maryland 21701

Contract No. DADA 17-72-C-2005

University of Maryland School of Medicine
Baltimore, Maryland 21201

Approved for public release; distribution unlimited

The findings in this report are not to be construed as
an official Department of the Army position unless so
designated by other authorized documents.



AD A 072400

DDC FILE COPY

79 08 06 095

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER 9
4. TITLE (and Subtitle) Hemoglobin Function in Stored Blood .		5. TYPE OF REPORT & PERIOD COVERED Annual Report - 1 February 1978 - 31 Jan 1979
7. AUTHOR(s) R. Ben/Dawson / M. D.		6. PERFORMING ORG. REPORT NUMBER CONTRACT OR ORDER NUMBER(s) DADA 17-72-C-2005
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of Maryland School of Medicine Baltimore, Maryland 21201		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 62772A 3S762772A814 00.042
11. CONTROLLING OFFICE NAME AND ADDRESS US Army Medical Research and Development Command Fort Detrick, Frederick, Maryland 21701		12. REPORT DATE 11 January 1979
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 12 7p		13. NUMBER OF PAGES 97
		15. SECURITY CLASS. (of this report) Unclassified
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Blood preservation, Red Cell Metabolism, Red Cell Function, 2,3-Diphosphoglycerate, Adenine, Inosine, Methylene Blue, Pyruvate, Dihydroxyacetone, Ascorbic Acid, D-ascorbate, Dehydroascorbate, Hexoses, Dissaccharides, Pentose.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This report presents ATP, 2,3-DPG, pH, and glucose levels of whole blood and packed cells stored in CPD-adenine with the following variables; pH, glucose concentrations, adenine, inosine, methylene blue, phosphate, pyruvate, dihydroxy- acetone, ascorbic acid, fructose, mannose, galactose, and ribose. Also, D- ascorbate, dehydroascorbate and SH inhibitors. 219400 km		

ANNUAL PROGRESS REPORT (31 Jan. 1979) ON

CONTRACT DADA-17-72-C-2005

HEMOGLOBIN FUNCTION IN STORED BLOOD

February 1, 1978 thru January 31, 1979

INTRODUCTION

During this contract year, work was continued on liquid preservation of stored blood for the purpose of maintaining 2,3-DPG levels for hemoglobin function and ATP levels for red cell post transfusion survival. Preservatives used were CPD (citrate-phosphate-dextrose), and CPDA-1. Other variables studied were concentrations of glucose, other 5 and 6 carbon sugars and three dissaccharides. Also, the metabolic regulators, methylene blue and ascorbate, D-ascorbate and dehydroascorbate, with sulfhydryl inhibitors. Other experiments used concentrations of phosphate, dehydroxyacetone, inosine and adenosine.

This report covers 11 months of contract work during which time 16 experiments were brought to completion. Some of these will not be covered in this report as data analysis is not complete on several. The 16 items given as the body of this report include the abstract of a previously submitted paper and abstracts submitted for publication on three of the papers whose entire manuscripts are given herein. Thus, in the 11 abstracts which are in this report, 7 give results not given previously or in this report; and, in addition there are 5 manuscripts submitted here for the first time.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist.	Avail and/or special
A	

RESULTS AND DISCUSSION

One of the important conclusions from our large, CPD-adenine packed cell study, Blood Preservation 26, is that high hematocrit packed cells (Hct 80-95%) require at least 175% of the glucose present in CPD for 35 days of storage. Also, for 42 days of storage, 200% of glucose seems to be indicated.

In BP 28, galactose and maltose were shown to be of some value in maintaining red cell 2,3-DPG and ATP, although not as good as fructose and dextrose.

In BP 33, phosphate, in increased concentrations was shown to have a beneficial effect on 2,3-DPG levels with ribose as well as inosine. This is a new finding and adds some impetus to the exploration of alternate sugars to glucose such as ribose.

Dihydroxyacetone (DHA) has been of some interest for 5 years, for its ability to maintain 2,3-DPG although its adverse effect on ATP has been a concern. In BP 34, increasing the phosphate concentration reduces the adverse effect of DHA on ATP. The extra phosphate, 10 mM, slightly lowers 2,3-DPG values; however, the levels are still maintained at approximately half normal for 4 weeks. The merit of this 2,3-DPG maintenance in a DHA-10 mM phosphate preservative, that maintains higher than control ATP values, should be further evaluated.

In another experiment, BP 35, DHA and D-ascorbate were evaluated together with phosphate in CPD-adenine. Again, ATP maintenance was improved with DHA and phosphate. However, DHA and D-ascorbate kept 2,3-DPG levels above normal for 6 weeks of storage. As in the experiments without ascorbate, phosphate improved ATP maintenance in DHA with ascorbate.

Our first experiment in the reevaluation of adenosine is reported here in abstract form, BP 40. 5 mM adenosine maintained 2,3-DPG at near normal for 21 days of storage and was clearly better than lower concentrations. However, ATP concentrations were depressed by adenosine. Of note was the higher glucose levels observed with this 5 mM preservative, possibly indicating glucose sparing as the ribose part of adenosine is utilized. BP 41 studied adenosine in a CPD-adenine

RESULTS AND DISCUSSION (CONT)

preservative with D-ascorbate. Although the adverse effect noted above adenosine on ATP is additive to the adverse effect of D-ascorbate, there is an important additive effect of adenosine and D-ascorbate on 2,3-DPG maintenance

In further studies on the ascorbic acid mechanisms of red cell maintenance of 2,3-DPG, BP 43, iodoacetate was shown to have a profound inhibitory effect on red cell metabolism, not protected by ascorbate or glutathione. In BP 44, dehydro-ascorbate, the oxidized form of the redox pair, was shown to have a slightly greater ability to maintain 2,3-DPG during blood storage than D-ascorbate, the unnatural isomer of the vitamin.

The ion exchange resin, Amberlite-45 with and without phosphate, was shown to stabilize blood preservative pH during storage in CPD, helping to maintain 2,3-DPG. Also, the phosphate loaded resin was presumed to release phosphate slowly during storage, thus facilitating maintenance of ATP.

Passage of oxygen through the plastic blood storage bag has been shown to raise the partial pressure of oxygen during liquid blood storage. It is supposed that a resultant increase in the fraction of oxyhemoglobin will release bound 2,3-DPG, thus inhibiting the formation of more 2,3-DPG during blood storage. The quantitative impact of this effect of gas permeable plastic blood bags is being assessed.

SUMMARY

With the licensing of CPDA-1 for 5 weeks of blood bank storage in Aug. 1978 and the adoption of this advance by the American National Red Cross in September, a major development in blood preservation research has been realized. The next major effort in this field; that is, maintaining normal 2,3-DPG levels for immediate oxygen transport upon transfusion now comes to the front among the unresolved issues in blood preservation and transfusion therapy. As this work has been going on, studies in animals and humans has shown improved function and survival when higher or normal 2,3-DPG blood is transfused, compared to the low 2,3-DPG blood, in certain control circumstances.

The present report summarizes the results of 11 abstracts and 5 manuscripts most of which are directly aimed at maintaining 2,3-DPG levels over 5-6 weeks of blood bank storage. The 5 manuscripts and 7 of the abstracts represent work not previously reported. Four of the abstracts are on work presented in part, previously. This laboratory expects to continue this work with some of the metabolic regulators and nutrients mentioned here and others as well.

DISTRIBUTION LIST

4 copies

HQDA (SGRD-SI)
Fort Detrick
Frederick, MD. 21701

12 copies

Defense Documentation Center (DDC)
ATTN: DDC- DDA
Cameron Station
Alexandria, Virginia 22314

1 copy

Dean
School of Medicine
Uniformed Services University of the
Health Sciences
4301 Jones Bridge Road
Bethesda, Maryland 20014

1 copy

Superintendent
Academy of Health Sciences, US Army
ATTN: AHS-COM
Fort Sam Houston, Texas 78234